

# MORE ON CYCLIC AMENABILITY OF THE LAU PRODUCT OF BANACH ALGEBRAS DEFINED BY A BANACH ALGEBRA MORPHISM

M. RAMEZANPOUR

Abstract. For two Banach algebras  $A$  and  $B$ , the  $T$ -Lau product  $A \times_T B$ , was recently introduced and studied for some bounded homomorphism  $T : B \rightarrow A$  with  $\|T\| \leq 1$ . Here, we give general necessary and sufficient conditions for  $A \times_T B$  to be (approximately) cyclic amenable. In particular, we extend some recent results on (approximate) cyclic amenability of direct product  $A \oplus B$  and  $T$ -Lau product  $A \times_T B$  and answer a question on cyclic amenability of  $A \times_T B$ .

## 1. Introduction and some Preliminaries

The notion of weak amenability for commutative Banach algebras was introduced and studied for the first time by Bade, Curtis and Dales [3]. Johnson [11] extended this concept to the non commutative case and showed that group algebras of all locally compact groups are weakly amenable. A Banach algebra  $A$  is called weakly amenable if every continuous derivation  $D : A \rightarrow A^*$  is inner. It is often useful to restrict one's attention to derivations  $D : A \rightarrow A^*$  satisfying the property  $D(a)(c) + D(c)(a) = 0$  for all  $a, c \in A$ . Such derivations are called cyclic. Clearly inner derivations are cyclic. A Banach algebra is called cyclic amenable if every continuous cyclic derivations  $D : A \rightarrow A^*$  is inner. This notion was presented by Gronbaek [9]. He investigated the hereditary properties of this concept, found some relations between cyclic amenability of a Banach algebra and the trace extension property of its ideals.

Ghahramani and Loy [7] introduced several approximate notions of amenability by requiring that all bounded derivations from a given Banach algebra  $A$  into certain Banach  $A$ -bimodules to be approximately inner. In the same paper and the subsequent one [8], the authors showed the distinction between each of these concepts and the corresponding classical notions and investigated properties of algebras in each of these new classes. Motivated by this notions, Esslamzadeh and Shojaei [6] defined the concept of approximate cyclic amenability for Banach algebras and investigated the hereditary properties for this new notion. Shojaei and

---

1991 *Mathematics Subject Classification*. Primary 46H05; Secondary 46H99.

*Key words and phrases*. Banach algebra, cyclic amenability,  $T$ -Lau product.

Bodaghi in [14, Theorem 2.3] showed that for Banach algebras  $A$  and  $B$ , if direct product  $A \oplus B$  with  $\ell^1$ -norm is approximately cyclic amenable, then so are  $A$  and  $B$ . They also showed that the converse for the case where  $A^2$  is dense in  $A$ .

On the other hand, for two Banach algebras  $A$  and  $B$  and a bounded homomorphism  $T : B \rightarrow A$  with  $\|T\| \leq 1$ , the  $T$ -Lau product  $A \times_T B$  is defined as the space  $A \times B$  equipped with the norm  $\|(a, b)\| = \|a\| + \|b\|$  and the multiplication

$$(a, b)(a', b') = (aa' + aT(b') + T(b)a', bb'),$$

for all  $a, a' \in A$  and  $b, b' \in B$ . This product was first introduced and studied by Bhatt and Dabhi in [4] for the case where  $A$  is commutative. Javanshiri and Nemati in [10] extended this product to the general Banach algebras and studied Arens regularity, amenability and  $n$ -weak amenability of  $A \times_T B$  in the general case; see also [2] and [5]. When  $T = 0$ , this multiplication is the usual coordinatewise product and so  $A \times_T B$  is in fact the direct product  $A \oplus B$ . Furthermore, let  $A$  be unital with the identity element  $e$  and let  $\theta : B \rightarrow \mathbb{C}$  be a non-zero multiplicative linear functional. Define  $T_\theta : B \rightarrow A$  as  $T_\theta(b) = \theta(b)e$ , for each  $b \in B$ . Then the above product with respect to  $T_\theta$  coincides with the product investigated in [12].

Bhatt and Dabhi in [4] showed that cyclic amenability of  $A \times_T B$  is stable with respect to  $T$ , for the case where  $A$  is commutative, but the proof contains a gap. In [1] Abtahi and Ghafarpanah fixed this gap and extended this result to an arbitrary Banach algebra  $A$ . Indeed they proved that if  $A \times_T B$  is cyclic amenable then both  $A$  and  $B$  are cyclic amenable. They also proved the converse for the case where both  $A$  and  $B$  have faithful dual spaces. But they left it open for all Banach algebras; [1, Question 3.5].

In the present paper, we give general necessary and sufficient conditions for  $A \times_T B$  to be (approximate) cyclic amenable. In particular we extend the recent results on (approximate) cyclic amenability of the direct product  $A \oplus B$ , [14], and the  $T$ -Lau product  $A \times_T B$  and answer Question 3.5 in [1] on cyclic amenability of  $A \times_T B$ .

## 2. Cyclic amenability

Let  $A$  be a Banach algebra, and  $X$  be a Banach  $A$ -bimodule. Then the dual space  $X^*$  of  $X$  becomes a dual Banach  $A$ -bimodule with the module actions defined by  $(fa)(x) = f(ax)$  and  $(af)(x) = f(xa)$ , for all  $a \in A, x \in X$  and  $f \in X^*$ . A derivation from  $A$  into  $X$  is a linear mapping  $D : A \rightarrow X$  satisfying

$$D(ac) = D(a)c + aD(c) \quad (a, c \in A).$$

If  $x \in X$  then  $d_x : A \rightarrow X$  defined by  $d_x(a) = ax - xa$  is a derivation which is called an inner derivation.

A derivation  $D : A \rightarrow A^*$  is said to be cyclic if  $D(a)(c) + D(c)(a) = 0$  for all  $a, c \in A$ . If every continuous cyclic derivation  $D : A \rightarrow A^*$  is inner then  $A$  is called cyclic amenable.

As remarked in [4], the dual space  $(A \times_T B)^*$  can be identified with the Banach space  $A^* \times B^*$  equipped with the norm  $\|(f, g)\| = \max\{\|f\|, \|g\|\}$  via

$$(f, g)(a, b) = f(a) + g(b),$$

where  $a \in A, f \in A^*, b \in B$  and  $g \in B^*$ . Moreover, a direct verification reveals that  $(A \times_T B)$ -module operations of  $(A \times_T B)^*$  are as follows.

$$\begin{aligned} (f, g)(a, b) &= (fa + fT(b), T^*(fa) + gb), \\ (a, b)(f, g) &= (af + T(b)f, T^*(af) + bg) \end{aligned}$$

for  $a \in A, b \in B, f \in A^*$  and  $g \in B^*$ .

To clarify the relation between cyclic amenability of  $A \times_T B$  and that of  $A$  and  $B$ , we need the next result which characterizes the continuous cyclic derivations on  $A \times_T B$ .

**Lemma 2.1.** *Suppose that  $A$  and  $B$  are Banach algebras and  $T : B \rightarrow A$  is a bounded (by one) homomorphism. Then every continuous cyclic derivation  $D : A \times_T B \rightarrow (A \times_T B)^*$  enjoys the presentation*

$$D(a, b) = (D_1(a) - S^*(b), D_2(b) + S(a))$$

for all  $a \in A$  and  $b \in B$ , where

- (a)  $D_1 : A \rightarrow A^*$  and  $D_2 : B \rightarrow B^*$  are continuous cyclic derivations.
- (b)  $S : A \rightarrow B^*$  is a bounded linear operator such that  $S(ac) = (T^* \circ D_1)(ac)$  and  $S(a)b = (T^* \circ D_1)(a)b$  for all  $a, c \in A$  and  $b \in B$ .

Moreover,  $D$  is inner if and only if  $S = T^* \circ D_1$  and both  $D_1$  and  $D_2$  are inner derivations.

Abtahi and Ghafarpanah in [1], proved that if  $A \times_T B$  is cyclic amenable then both  $A$  and  $B$  are cyclic amenable. They also proved the converse for the case where both  $A$  and  $B$  have faithful dual spaces, but left it as an open question for all Banach algebras; [1, Question 3.5]. An improvement of this result has been also obtained in [13]. Indeed, the converse has been proved for the case where  $A^2$  is dense in  $A$ ; [13, Theorem 2.6]. We should remark that a Banach algebra  $A$  has left (right) faithful dual space just when  $A^2$  is dense in  $A$ .

Here we give general necessary and sufficient conditions for  $A \times_T B$  to be cyclic amenable. This result improves [13, Theorem 2.6] and answers also Question 3.5 in [1].

**Theorem 2.2.** *Suppose that  $A$  and  $B$  are Banach algebras and  $T : B \rightarrow A$  is a bounded (by one) homomorphism. Then  $A \times_T B$  is cyclic amenable if and only if the following statements hold.*

- (1)  $A$  and  $B$  are cyclic amenable.
- (2)  $A^2$  is dense in  $A$  or  $B^2$  is dense in  $B$ .

### 3. Approximate cyclic amenability

Recall from [7] that a derivation  $D : A \rightarrow X$  is called approximately inner if there exists a net  $\{x_\alpha\} \subseteq X$  such that  $D = \lim_\alpha d_{x_\alpha}$  in the strong operator topology. A Banach algebra  $A$  is called approximately cyclic amenable, if every continuous cyclic derivation  $D : A \rightarrow A^*$  is approximately inner. The concepts of approximate cyclic amenability was introduced and studied in [6]; see also [14].

The next result characterizes approximately inner derivations on  $A \times_T B$ .

**Proposition 3.1.** *Suppose that  $A$  and  $B$  are Banach algebras and  $T : B \rightarrow A$  is a bounded (by one) homomorphism. Let  $D : A \times_T B \rightarrow (A \times_T B)^*$  be a continuous cyclic derivation given by  $D(a, b) = (D_1(a) - S^*(b), D_2(b) + S(a))$  for all  $a \in A$  and  $b \in B$ . Then  $D$  is approximately inner if and only if  $S = T^* \circ D_1$  and both  $D_1$  and  $D_2$  are approximately inner.*

Applying Proposition 3.1 and using a similar argument used in the proof of Theorem 2.2 we can prove the next theorem. This result improves [13, Theorem 2.3].

**Theorem 3.2.** *Suppose that  $A$  and  $B$  are Banach algebras and  $T : B \rightarrow A$  is a bounded (by one) homomorphism. Then  $A \times_T B$  is approximately cyclic amenable if and only if the following statements hold.*

- (1)  $A$  and  $B$  are approximately cyclic amenable.
- (2)  $A^2$  is dense in  $A$  or  $B^2$  is dense in  $B$ .

Shojaee and Bodaghi in [14, Theorem 2.3] showed that for Banach algebras  $A$  and  $B$ , if direct product  $A \oplus B$  is approximately cyclic amenable, then so are  $A$  and  $B$ . They also showed that the converse for the case where  $A^2$  is dense in  $A$ . Applying Theorems 2.2 and 3.2 for  $T = 0$ , we get the next result which extends [14, Theorem 2.3].

**Corollary 3.3.** *Let  $A$  and  $B$  be Banach algebras. Then the direct product  $A \oplus B$  is (approximately) cyclic amenable if and only if the following statements hold.*

- (1)  $A$  and  $B$  are (approximately) cyclic amenable.
- (2)  $A^2$  is dense in  $A$  or  $B^2$  is dense in  $B$ .

Let  $A$  be unital and  $\theta : B \rightarrow \mathbb{C}$  be a non-zero multiplicative linear functional. Define  $T_\theta(b) := \theta(b)e$ . Then  $A \times_{T_\theta} B$  is the  $\theta$ -Lau product  $A \times_\theta B$ , [12]. As a consequence of Theorems 2.2 and 3.2, we have the next result.

**Corollary 3.4.** *Let  $A$  be unital and  $\theta$  be a non-zero multiplicative linear functional on  $B$ . Then  $A \times_\theta B$  is (approximately) cyclic amenable if and only if both  $A$  and  $B$  are (approximately) cyclic amenable.*

## References

- [1] Abtahi, F.—Ghafarpanah, A.: *A note on cyclic amenability of the Lau product of Banach algebras defined by a Banach algebra morphism*, Bull. Aust. Math. Soc., **92** no. 2 (2015), 282–289.
- [2] Abtahi, F.—Ghafarpanah, A.—Rejali, A.: *Biprojectivity and biflatness of Lau product of Banach algebras defined by a Banach algebra morphism*, Bull. Aust. Math. Soc., **91** no. 1 (2015), 134–144.
- [3] Bade, G. W.—Curtis, P. C.—Dales, H. G.: *Amenability and weak amenability for Beurling and Lipschitz algebras*, Proc. London Math. Soc., **55** no. 2 (1987), 359–377.
- [4] Bhatt, S. J.—Dabhi, P. A.: *Arens regularity and amenability of Lau product of Banach algebras defined by a Banach algebra morphism*, Bull. Aust. Math. Soc., **87** (2013), 195–206.
- [5] Dabhi, P. A.—Jabbari, A.—Haghnejad Azar, K.: *Some notes on amenability and weak amenability of Lau product of Banach algebras defined by a Banach algebra morphism*, Acta Math. Sin. (Engl. Ser.), **31** no. 9 (2015), 1461–1474.
- [6] Esslamzadeh, G. H.—Shojaee, B.: *Approximate weak amenability of Banach algebras*, Bull. Belg. Math. Soc. Simon Stevin, **18** no. 3 (2011), 415–429.
- [7] Ghahramani, F.—Loy, R. J.: *Generalized notions of amenability*, J. Funct. Anal., **208** no. 1 (2004), 229–260.
- [8] Ghahramani, F.—Loy, R. J.—Zhang Y.: *Generalized notions of amenability. II*, J. Funct. Anal., **254** no. 7 (2008), 1776–1810.
- [9] Grønbæk, N.: *Weak and cyclic amenability for noncommutative Banach algebras*, Proc. Edinburgh Math. Soc., **35** no. 2 (1992), 315–328.
- [10] Javanshiri, H.—Nemati, M.: *On a certain product of Banach algebras and some of its properties*, Proc. Rom. Acad. Ser. A, **15** no. 3 (2014), 219–227.
- [11] Johnson, B. E.: *Derivations from  $L^1(G)$  into  $L^1(G)$  and  $L^\infty(G)$* , Lecture Notes in Math., 1359, 191–198, Springer, Berlin, 1988.
- [12] Sangani Monfared, M.: *On certain products of Banach algebras with applications to harmonic analysis*, Studia Math., **178** (3) (2007), 277–294.
- [13] Nemati, M.—Javanshiri, H.: *Some cohomological notions on  $A \times_T B$* , ArXiv:1509.00894v2, (2015).
- [14] Shojaee, B.—Bodaghi, A.: *A generalization of cyclic amenability of Banach algebras*, Math. Slovaca, **65** (3) (2015), 633–644.

School of Mathematics and Computer Science, Damghan University, P. O. Box 36716, Damghan  
41167, Iran.

*E-mail address:* `ramezanpour@du.ac.ir`